

**A SURVEY OF WRITING TECHNIQUES,
AUTHORSHIP, TENSES, VOICES,
AND LENGTH OF OPENING SENTENCES
OF MEDICAL RESEARCH ARTICLES**

Phan Banpho

Chulalongkorn University Language Institute

Abstract

A large number of genre analyses have investigated schematic units or “moves” of many parts of medical research articles. The present analysis deals with the inner element of the first move, the introduction section. It explores the techniques for writing opening sentences, and investigates authorship, tenses, voices, and sentence length. The subjects were 167 opening sentences from two world journals in medical science, the *Annals of Internal Medicine* and *The Lancet*, which represent American English and British English, respectively. A concordancing computer program, Microsoft Word, and the SPSS (v.11.5) were used. The findings of this study suggest that most opening sentences, which comprise an average of about 22 words, give facts in the present simple active by citation of previous findings.

Introduction

The opening or first sentence of the introduction section of a research article is crucial, partly because it should capture the readers’ attention (McElroy, 1997, p. 40) as well as introduce or give a lead into the research topic under investigation. Article

writers, especially new entrants into the academic discourse community, often find it difficult and time-consuming to write this first sentence. Several writing techniques have been recommended. McElroy (1997, pp. 40-41) suggests the following writing techniques, or rhetorical functions, to catch the reader's attention: giving facts, statistics, a definition, and a general statement. Swales (1984, p. 81) has recommended that the opening sentence should attempt to establish that the field of research is useful, relevant, or important in some way or ways. Also, in his revised model for article introduction, CARS (Create a Research Space), he has established that the first move of the introduction should claim that the area of the study is an important one, implying that the topic under study is also crucial (Swales, 1990). Brink and Wood (1994, p. 239) have proposed that the first sentence in the introduction should set the general tone and direction for the subject matter. Unfortunately, there is no empirical data about rhetorical functions, or techniques, for writing the opening sentence of medical research articles. In addition, no study has been undertaken to investigate whether the opening sentence is written by the authors of the article themselves or is a citation of previous findings.

The use of tenses and voices is another aspect of scientific genre that has been investigated extensively. Wingard (1981) has found that in a corpus of six medical texts, the most frequent form is the simple present active (including *is/are*), accounting for 28% of cases, followed by the simple past active (22%). The simple present passive accounts for 11%, while the simple past passive 18%. Barber (1985, p. 8) has analyzed the frequency of different verb tenses and voices, some of which are as follows: the present simple active (64%), present simple passive (25%), present perfect passive (1.7%), present perfect active (1.4%), past simple active (1.2%), and past simple passive (1.2%). Again, these studies do not focus primarily on the opening sentence of medical research articles.

Sentence length is another important issue that worries many authors because it may affect readability, influence the credibility of the results and, worst of all, distract the reader. A very short sentence may be inadequate, whereas an excessively

long one may fail to hold the readers' attention. According to Barber (1985), the average sentence length in scientific prose is 27.6 words. However, there is no data regarding the average length of the opening sentences in medical research articles.

Although a number of studies on medical research articles have been carried out, there is no study that focuses on the first inner element of the introduction section, the opening sentence. Therefore, the purposes of this study were 1) to explore techniques, or rhetorical functions, for generating opening sentences, 2) to investigate whether these first sentences were written by the authors or were just a citation of previous findings, and 3) to examine the tense, voice and length of these opening sentences.

Methods

Subjects

There is an enormous number of professional journals in the medical sciences. Some are released weekly, some biweekly, while others come out once a month. Furthermore, some are written in American English, others in British English. This research studied two medical journals—the *Annals of Internal Medicine*, which is an American professional journal that is published biweekly, and *The Lancet*, a British journal published weekly. The main reasons for choosing these two journals are that neither focuses on a particular medical discipline and that they are two leading world journals. These journals present several kinds of articles such as general articles, research articles (or original articles), and case reports, but only research articles were studied in this study.

The initial population of this study consisted of 292 opening sentences, 72 from the *Annals of Internal Medicine* and 220 from *The Lancet*. These articles were published in 2002. However, to avoid having an overwhelming number of opening sentences from *The Lancet*, only the first and the third issues of this journal were assessed. The actual subjects of this study became 167 sentences—95 from *The Lancet* and 72 from the *Annals of Internal Medicine*.

Instruments

The research instruments used were coding forms and three computer programs: Microsoft Word, a concordancing program (or concordancer), and the SPSS (v.11.5).

Procedures

Data collection

For data collection, the first page of 167 articles was photocopied. For reference and ease of recognition, each opening sentence was numbered at the beginning and end of the sentence as in the following example: #1011 *Chronic obstructive pulmonary disease (COPD) is one of the most important problems world-wide (1,2) (#1011 Feb 19/1)*. The opening sentences from the two sources—*The Lancet* and the *Annals of Internal Medicine*—were typed and saved as Corpus 1 and Corpus 2, respectively.

Table 1: Month, dates and numbers of articles

| Months | Annals | | | Lancet | | | Total |
|-----------|-----------|---------|--------------|-----------|---------|--------------|------------|
| | Date | (Number | of articles) | Date | (Number | of articles) | |
| January | 1, 15 | (6) | | 5, 19 | (8) | | 14 |
| February | 5, 19 | (6) | | 2, 16 | (8) | | 14 |
| March | 5, 19 | (6) | | 2, 16 | (8) | | 14 |
| April | 2, 16 | (6) | | 6, 20 | (8) | | 14 |
| May | 7, 21 | (6) | | 4, 18 | (8) | | 14 |
| June | 4, 18 | (6) | | 1, 15 | (8) | | 14 |
| July | 2, 16 | (6) | | 6, 20 | (8) | | 14 |
| August | 6, 20 | (6) | | 3, 17 | (8) | | 14 |
| September | 3, 17 | (6) | | 7, 21 | (8) | | 14 |
| October | 1, 15 | (6) | | 5, 19 | (8) | | 14 |
| November | 5, 19 | (6) | | 2, 16 | (8) | | 14 |
| December | 3, 17 | (6) | | 7, 21 | (7) | | 13 |
| | 72 | | | 95 | | | 167 |

Note: January 1, 15 (6) refers to January 1, 2002, January 15, 2002 and (6) refers to 6 articles.

Writing techniques
Table 2: Wordlist (adjectives)

| Word | Frequency |
|------------------------|-----------|
| 1. most | 10 |
| 2. acute | 9 |
| 3. common | 7 |
| 4. million | 6 |
| 5. major | 5 |
| 6. primary | 5 |
| 7. chronic | 4 |
| 8. important | 4 |
| 9. public | 4 |
| 10. <i>high</i> | 3 |
| 11. <i>long</i> | 3 |
| 12. <i>severe</i> | 3 |
| 13. <i>short</i> | 3 |
| 14. <i>substantial</i> | 3 |
| 15. <i>serious</i> | 2 |
| Total | 71 |

To assess the writing techniques, McElroy's suggestion and Swales' recommendation were initially used to form a tentative rhetorical classification. The initial rhetorical functions or writing techniques derived from McElroy and Swales were "*indicating importance/usefulness/relevance*," "*giving facts*," "*expressing comments/observations*," "*giving definitions*," and "*describing situations*." In this study, "*facts*" includes both general facts, generalizations, and findings cited from previous research findings. Regarding importance, the opening sentence that would be considered as expressing the rhetorical function "*indicating importance*" must have at least one essential characteristic, containing an adjective, a noun, or an adverb that signals importance, such as *important*, *mortality*, and *highly*.

To have a more precise sketch of the writing techniques, Corpus 1 and Corpus 2 were loaded into the concordancer to display a wordlist and frequency. Then, some potential words with high frequency, particularly those that might lead to

identification of the actual writing techniques used in the study samples, were selected. To avoid having a complicated list of concordances (or a list of words from a text or group of texts, displayed straight down from the top to bottom of the page as in Figure 1), potential words were divided into groups, such as *adjectives*, *nouns*, *verb to be*, and *modals*. When individual groups of potential words were identified, some low frequency words belonging to the same group as the high frequency ones were added to the list. Table 2 displays potential adjectives that might be an important element of "*indicating importance*." The italicized words in Table 2 are words with low frequency added to the initial list. Finally, potential key words were used as key words to generate concordances as shown in Figure 1.

Figure 1: Concordances of adjectives from Corpus 1 and Corpus 2

| | |
|---|---|
| #1018 Feb2 /2)*#1019 | Asthma is the most common chronic disease of childbirth |
| (#1068 May 4 / 1)#1069 | Most leg ulcers are venous.1 (#1069 M |
| May 4 / 3)*#1071 | Measles is the most contagious disease known to man. |
| 128 Ovarian carcinoma is the sixth most | common cancer of women worldwide and ac |
| mangliomas (strawberry naevi) are the | most common soft tissue tumours of infan |
| (#1130 Aug 17 / 3)#1139 | The most appropriate revascularisation strat |
| .1 (#1148 Sept. 21 / 2)#1149 | Most reviews of risk factors for asthma |
| nhibitors and aspirin are two of the | most widely used therapeutic agents in c |
| ated low back pain can be one of the | most challenging disorders seen in prime |
| (#1180 Nov 16 / 2)#1181 | One of the most important milestones in HIV-1 rese |
| 05 Colorectal cancer is the third | most common cause of cancer-related deat |
| monary disease (COPD) is one of the | most important problems world-wide (1,2) |
| ansferase (ALT) concentration is the | most commonly used variable for assessm |
| 055 Congestive heart failure is the | most frequent cause of hospitalization : |
| ong U.S. women, breast cancer is the | most common type of cancer and is the s |
| he common cold is one of the world's | most prevalent illness. (#2070 Dec 1' |
| ccurs spontaneously in patients with | acute coronary syndromes or during a pe |
| on lowers mortality in patients with | acute myocardial infarction. (#1017 |
| tcome.1-2 (#1025 Feb16 /2)*#1026 | Acute liver failure resulting from seve |
| ment of multiple organ systems, with | acute renal failure, haemodynamic instal |
| ressive agents for the prevention of | acute rejection in renal-allograft reci |
| reperfusion therapy in patients with | acute myocardial infarction depends on |
| nd major cardiovascular events after | acute myocardial infarction. 1-3 (#10 |
| n Africa. (#1114 Jul 20 /3)*#1115 | Acute bacterial meningitis causes many |
| 39 Sept. 7 / 1)#1140 | Patients with acute myocardial infarction and subst |
| cies, including hypertensive crisis, | acute myocardial infarction, and ventr |
| 1)#2038 | Emergent reperfusion for |
| se of cardiac catheterization after an | acute myocardial infarction (AMI) can b |
| illness. (#2070 Dec 17 / 1)#2071 | Acute myocardial infarction are mixed. |
| Feb2 /2)*#1019 | Asthma is the most common chronic disease of childbirth; a |
| 0 /1)* #1113 | Heart failure is a common and growing problem associated w |
| Ovarian carcinoma is the sixth most | common cancer of women worldwide and ac |
| omas (strawberry naevi) are the most | common soft tissue tumours of infancy, |
| reatment of choice for patients with | common bile-duct stones.1-3 |
| sical inactivity, and obesity -- are | common to both stroke and coronary hear |
| hepatitis B Virus (HBV) and HIV-1 is | common, because of shared modes of tra |
| Colorectal cancer is the third most | common cause of cancer-related death in |
| / 1)#2020 | Nocturnal hypoglycemia is common in patients with type 1 diabetes |
| emity peripheral arterial disease is | common, both in community studies of un |
| 2)#2057 | Venous thromboembolism is common after orthopedic surgery. (#2057 |
| .S. women, breast cancer is the most | common type of cancer and is the second |
| (1-3). (#2069 Dec 3 / 3)#2070 | The common cold is one of the world's most |
| (#1020 Feb2 /4)*#1024 | About 5 million people are imprisoned worldwide |
| re imprisoned worldwide, including 2 | million in the USA and 70 000 in the UK |
|)*#1093 | More than 70% of the 36.1 million HIV-1-infected individuals we |
| Onchocera volvulus infects about 18 | million people, *99% of whom live in su |
| / 2)**#1122 | There are up to half a million new cases of visceral leishmani |
| At the end of 2000, an estimated 35 | million people had been displaced by * |
| od pressure, which affects nearly 50 | million Americans, is a serious public |
| hronic disease, currently affects 16 | million people in the United States, c |
| ductase reduce the risk of death and | major cardiovascular events after acute |
| function in young children may have | major implications for our understandin |
| #1095 | Type 2 diabetes mellitus is a major health problem associated with ex |
| monary hypertension, which can be a | major cause of morbidity and mortality. |

The concordances in Figure 1 indicate mixed rhetorical functions, particularly "giving fact," "expressing a combination of facts and importance," and "indicating importance."

After this first try, potential nouns were used as key words to generate concordances. Examples of nouns with high frequency and those potential words are shown in Figure 2. The

concordances in this figure reveal "giving fact," "indicating importance," and "using a combination of fact and importance." Examples of nouns that might suggest the writing technique "indicating importance", were mortality, morbidity, death(s), disease(s), risk, cause(s), severity, life, and cost(s).

Figure 2: Concordances of nouns from Corpus 1 and Corpus 2

tanecous coronary intervention lowers mortality in patients with acute myocardial infarction. (#1040 Mar 16 / 1)*#1041 T
 th the aim to decrease breast cancer mortality. (#1040 Mar 16 / 1)*#1041 T
 an be a major cause of morbidity and mortality.1 (#1148 Sept 21 / 2)#1
 / 4)#1163 Despite reductions in the mortality of patients with ST-elevation
 ast several decades, the short-term mortality rates remains high in two subc
 llution is associated with increased mortality from respiratory and cardiovas
 blood pressure and vascular disease mortality in particular epidemiological
 blood pressure levels, for which the mortality rates are relatively new. (#11
 ry bypass continues to be a cause of morbidity associated with cardiac surger
 alth problem associated with excess *morbidity and results in substantial hear
 ny deaths and substantial long-term *morbidity throughout the world.1 (#
 nsion, which can be a major cause of morbidity and mortality.1 (#1148 s
 n result in myocardial infarction or death. (#1009 Jan 19 / 1)* #1010
 MC-CoA) reductase reduce the risk of death and major cardiovascular events af
 at reducing the 30-day composite of death, myocardial infarction, and urgent
 ute bacterial meningitis causes many deaths and substantial long-term *morbid
 accounts for the greatest number of deaths from gynaecological malignancy in
 particulate air pollution and daily deaths.1-3 (#1166 Oct. 19 / 4)#1171
 1 aortic aneurysms caused about 6800 deaths in England and Wales in the year
 ted with resorption of embryo, fetal deaths, and potentially lethal malformat
 openic purpura (AITP) are at risk of life-threatening bleeding and therefore
 transplantation (LDLT) has become a life-saving option for some patients in
 ation between lung function in early life and lung disease in adulthood.1 (#
 in the pathogenesis of peptic-ulcer disease is controversial, because studie
 en in the UK have moderate or severe disease.3 (#1011 Jan 19 / 3)*#1012 T
 9 Asthma is the most common chronic disease of childbirth; asthma prevalence
 tself is the culprit, or whether the disease is caused by a different compone
 arthritis is a chronic progressive disease associated with systemic inflam
 is a chronic inflammatory rheumatic disease with a prevalence of 0.5-1.9% fc
 s. (#1063 Apr 20 / 4)*#1068 Crohn's disease is a chronic inflammatory disord
 071 Measles is the most contagious disease known to man. (#1071 May 4
 erstanding of respiratory health and disease, especially in view of the assoc
 lung function in early life and lung disease in adulthood.1 (#1087 Jun 1 /
 Jun 1 / 4)*#1092 Peripheral artery disease affects up to 15% of adults over
 4 Patients with advanced Hodgkin's disease have an excellent outlook if tre
 relationship between coronary heart disease risk about blood LDL cholesterol
 developing another primary malignant disease is an issue of increased importa
 y Henoch-Sch nlein purpura, Kawasaki disease. (#1164 Oct. 19 / 2)#1165 Resu
 from respiratory and cardiovascular disease and from lung cancer. (#1165 Oc
 is an important cause of diarrhoeal disease in children and adults in develc
 on to both stroke and coronary heart disease, discordant patterns of these tw
 between blood pressure and vascular disease mortality in particular epidemic
 ons of magnitude and distribution of diseases and injuries, and their causes
 se, discordant patterns of these two diseases in relation to geography, ethni
 because studies examining these two risk factors in this order have had conf
 rombocytopenic purpura (AITP) are at risk of life-threatening bleeding and th
 Moderate hyperhomocysteinaemia is a risk factor for venous *thromboembolism.
 me *A (HMG-CoA) reductase reduce the risk of death and major cardiovascular e
 placement surgery are in the highest risk category for postoperative venous T
 nship between coronary heart disease risk about blood LDL cholesterol concent
 tration is not associated with lower risk.1-5 (#1104 Jul 6 / 1)*#1105 LDL
 contentious for patients at moderate risk. (#1139 Sept. 7 / 1)#1140 Patien
 ith breast cancer have improved, the risk of developing another primary malign
 Sept. 21 / 2)#1149 Most reviews of risk factors for asthma recommend extend
 that although many of the well-known risk factors -- such as hypertension, sm

After some of the writing techniques were extracted from the concordances, individual opening sentences were assessed manually.

Authorship

To identify authorship, every opening sentence was carefully examined to see if it was cited from one or more previous findings or written by the authors. For *The Lancet*, each sentence was examined for a raised number or numbers indicating footnoting, such as the following:

Patients who have had a stroke often have long-term difficulties with walking and other daily activities such as getting out of a chair and climbing stairs.¹

For the *Annals of Internal Medicine*, the opening sentences were examined for the number or numbers in parentheses, such as the following:

Colorectal cancer is the third most common cause of cancer-related death in men and women (1).

The frequencies were tabulated in coding forms and the percentage of the sentences composed using citation and those written by the authors were calculated.

Tenses

In this study, only the tense of the main clause of the opening sentence was examined. Tenses were initially assessed using concordances in Figure 1, together with concordances derived from the following key words: *is, are, has, have, was, were,* and *had* because these words together with the verbs in the sentence could indicate tenses. After obtaining a sketch of tenses used in the opening sentences, the tense in every opening sentence was examined manually.

Voice

Like tenses, voices were initially assessed by using concordances generated by such key words as *is, are, was, were,*

has been, have been, and had been. When a sketch of voices was obtained, the voice of each opening sentence was carefully examined.

Length

The number of words in each opening sentence was counted using the *Word Count* function from the menu “Tools” of the Microsoft Word software program. Finally, the maximum and minimum numbers of words were identified and the average number of words in the opening sentences was calculated.

Statistics

The data obtained was processed to determine the percentages, means, and standard deviations. Both Chi-square and Student t-tests were used to test differences between the two groups of subjects.

Results

This study investigated five aspects of the opening sentence of the introduction of medical research articles. It explored the techniques used for writing the opening sentence of research articles, by whom it was written—the authors or citation of previous works, tenses, voices, and finally it examined the length of the opening sentences. It was found that the three most common writing techniques were “*giving facts*” (53.89%), “*using a combination of facts and importance*” (24.55%), and “*indicating importance*” (14.37%) (Table 3).

Table 3: Techniques used in writing opening sentences

| Technique | Annals | Lancet | Annals & Lancet | Chi-square test |
|----------------------|-----------------------|-----------------------|-----------------------|-------------------|
| | (N = 72) | (N = 95) | (N = 167) | |
| 1. Fact | 42 (58.33%) | 48 (50.53%) | 90 (53.89%) | $\chi^2 = 5.04^*$ |
| 2. Importance | 14 (19.44%) | 10 (10.53%) | 24 (14.37%) | |
| 3. Fact & Importance | 11 (15.28%) | 30 (31.58%) | 41 (24.55%) | |
| 4. Others | 5 (6.94%) | 7 (9.72 %) | 12 (7.19 %) | |
| Total | 72 | 95 | 167 | |

* $p = .05$

However, “using a combination of fact and importance” was used more often in *The Lancet* (31.58%) than in the *Annals of Internal Medicine* (15.28%). The differences of the first three techniques were statistically significant ($p = .05$). As regards importance, it was found that many adjectives, such as *important*, *major*, *main*, *primary*, *prevalent*, *severe*, and *significant* were used. In addition, certain nouns were also used to indicate importance. Examples of these nouns were *death*, *mortality*, *morbidity*, *severity*, *factors*, *cause*, and *hospitalization* because these nouns express important things about our health and our lives, such as loss of life, intensity of sickness, and indispensable medical treatment. However, in some contexts they might suggest fact rather than importance. For instance, one of the following sentences expresses fact while the other importance:

1. The mortality rate of this disease is 1/100,000. (*Fact*)
2. This disease causes both morbidity and mortality. (*Importance*)

Table 4: Authorship of information for opening sentences

| Authorship | Annals (N = 72) | Lancet (N = 95) | Annals & Lancet (N = 167) | Chi-square Test |
|---|---------------------------|---------------------------|---|----------------------------|
| 1. Original (authors) | 25 (34.72 %) | 41 (43.16%) | 66 (39.52 %) | $\chi^2 = 1.22^*$ |
| 2. Citation (previous works) | 47 (65.28%) | 54 (56.84 %) | 101 (60.48 %) | |
| Total | 72 | 95 | 167 | |

* $p = .05$

Regarding authorship, it was found that most opening sentences of both the *Annals of Internal Medicine* and *The Lancet* were composed by citation of one or more previous findings (60.48%). The *Annals of Internal Medicine* used more citation than *The Lancet*, 65.28% and 56.84%, respectively (Table 4). However, the difference was not statistically significant ($p = .05$).

As regards tense, this study found that more than half (70.06%) of the opening sentences in both medical journals were expressed in the present simple tense. *The Lancet* used more present simple tense (76.84%) than the *Annals of Internal Medicine* (61.11%). The difference was statistically significant ($p = .05$) (Table 5).

Table 5: Tense in opening sentences

| Tenses | Annals (N = 72) | Lancet (N = 95) | Annals & Lancet (N = 167) | Chi-square test |
|------------------------------|---------------------------|---------------------------|---|----------------------------|
| 1. Present simple | 44 (61.11 %) | 73 (76.84%) | 117 (70.06%) | $\chi^2 = 5.04^*$ |
| 2. Present perfect | 17 (23.61%) | 12 (12.63 %) | 29 (17.37%) | |
| 3. Others | 11 (15.28%) | 10 (10.53%) | 21 (12.57%) | |
| Total | 72 | 95 | 167 | |

* $p = .05$

As for voice, it was found that more than three-fourths of the opening sentences in both journals were written in active voice (77.84%), whereas only 22.16% were in passive voice. When choice of voice usage was compared, the data obtained revealed that active voice was used more often than passive voice in both journals, 77.84% and 22.16%, respectively. *The Lancet* used more active voice than the *Annals of Internal Medicine*, 83.16% and 70.83%, respectively. However, the difference was not statistically significant ($p = .05$).

Table 6: Voice in opening sentences

| Voices | Annals (N = 72) | Lancet (N = 95) | Annals & Lancet (N = 167) | Chi-square test |
|---------------------|------------------------|------------------------|---------------------------------|--------------------|
| 1. Active voice | 51 (70.83 %) | 79 (83.16%) | 130 (77.84%) | $\chi^2 = 3.61^*$ |
| 2. Passive voice | 21 (29.17%) | 16 (16.84 %) | 37 (22.16%) | |
| Total | 72 | 95 | 167 | |

* $p = .05$

Finally, this study found that the average length of the opening sentences in both medical journals was 21.89 words, with a maximum of 70 and a minimum of 5 words. However, there was no statistically significant difference between the lengths of the opening sentences in both journals ($p = .05$).

Table 7: Length of opening sentences

| Length | Annals | Lancet | Annals & Lancet | t-test |
|---------|-----------------------|----------------------|----------------------|-----------|
| | (N = 72) | (N = 95) | (N = 167) | |
| Average | 21.79 (SD = 10.37) | 21.96 (SD = 9.52) | 21.89 (SD = 9.89) | |
| Maximum | 70 | 50 | 70 | t = .107* |
| Minimum | 6 | 5 | 5 | |
| | 72 | 95 | 167 | |

* $p = .05$

Discussion

The purposes of this study were to examine the techniques used in writing the opening sentence of the introduction section of medical research articles, and to investigate the authorship, tenses, voices, and length of these initial sentences. It was found that more than half of the opening sentences (53.89%) express facts, present a combination of facts and importance (24.55%), or indicate importance (14.37%). This finding confirms the technique of “*giving facts*” suggested by McElroy (1997), and “*indicating importance*” recommended by Swales (1987). However, it should be noted that “*facts*” refers to both general facts and facts reported from research findings. This finding, therefore, is also supported by the second finding, that more than half of the opening sentences (60.48%) were citations of previous findings.

It is interesting that the substance or content of the opening sentences was not generally initiated by the authors. Rather, it was a citation of previous research findings. This finding probably makes sense in that, by nature, research findings are empirical and as such rhetorically convincing. In addition, writing of the opening sentence is not an easy task. Also, it may be more convenient to write from information on hand than from scratch. Therefore, citing previous works predominates in the authors' writing.

This study also found that the majority of the opening sentences were expressed in present simple tense (70.06%),

followed by the present perfect tense (17.37%), and other tenses (12.57%), which included the past simple tense and the present progressive tense. This finding corresponds with the finding that “*giving facts*” predominates. This may be because facts are usually expressed in present simple tense, particularly natural facts or generalizations, which are by nature timeless. Oster (1981) in her research on the use of tenses in reporting past literature has proposed that the present tense is used to refer to past literature and to refer to quantitative results of past literature that are supportive or non-relevant, whereas the present perfect is used to make general statements about past literature. Malcolm (1987) in a study on tense usage in the *Journal of Pediatrics* has hypothesized that generalizations are mainly expressed in the present tense, references to a specific study in the past tense, and references to an area of inquiry in the present perfect tense. Malcolm has concluded that the choice of tense is ultimately a rhetorical one; the authors are free to choose the time location that best suits their purposes. Moreover, Biber et al. (1999, pp. 456-458) have stated that present tense verbs are somewhat more common than past tense verbs in academic prose. Most importantly, Biber et al. have also asserted that the present tense can be used to refer to events in the past, to present states, to present habitual behavior, or to introduce future events. In fact, the present tense is used to imply a lack of time restriction. McCarthy (1991, p. 62) has stated that the tenses and aspects do not seem to depend on time as much as to issues such as the sender's purpose, focus on different elements of the message, and the projection of a shared framework within which the receiver will understand the message.

As regards voice, this study found that the active voice was used more than the passive voice, 77.84% and 22.16%, respectively. This finding is consistent with that of Wingard (1981) who found greater use of active verbs than of passive verbs. However, it should be noted that Wingard studied verb forms and functions in a corpus of four medical papers, and not the opening sentences. Moreover, as regards tense and voice, Tarone et al. (1981) have concluded that the writer's communicative purpose governs choices at the grammatical and lexical levels.

Finally, this study found that the average length of the opening sentences in both the American and English journals was about 21.89 words, with a maximum of 70 and a minimum of 5 words. This finding is inconsistent with that of Barber (1985), who reported that the average sentence length was 27.6 words. However, Barber noted that his investigation was merely preliminary and beside this it dealt with astronomy, not medicine. Moreover, although the present study found that the average length of the opening sentences was 21.89 words, there is no rule governing sentence length in other inner elements of the first schematic unit or move as other moves in medical research articles. However, an excessively long sentence is likely to be uninteresting as well as boring, while an extremely short one could be inadequate.

Implications for second and foreign language teaching

This study suggests some implications for second and foreign language teaching as follows:

1. Writing the opening sentence is difficult. To alleviate this problem, several practical techniques should be introduced to students, including the three most common techniques found in this study: “*giving facts*,” “*using a combination of facts and importance*,” and “*indicating importance*.” Also, the teacher should introduce citations to facilitate student writing of the opening sentence.

2. Criteria for choices of tenses should be taught in a succinct manner, particularly in reporting on past literature. For instance, the following three sentences have different meanings:

1. Robinson (1973) *found* that diabetes was a costly disease.
2. Robinson (1973) *finds* that diabetes is a costly disease.
3. Robinson (1973) *has found* that diabetes is a costly disease.

The first example refers to a specific study in the past. The second indicates a generalization that diabetes is a disease that costs a lot of money, regardless of time. The third example is to

claim generality about past literature—that diabetes is still an expensive disease from past to present.

3. In teaching tenses, the teacher should make it clear that the choice of tense *does not depend on time location only*, but also on the rhetorical function (illocutionary force), or the communicative function of the sentence as shown in the above examples.

4. This study found that the majority of first sentences in both *The Lancet* and *Annals of Internal Medicine* were composed by citation of previous works (60.48%). This indicates that citation of previous works is a practical and useful writing technique because both the content and structure of the sentence are already available. In addition, since the opening sentence is very important, many novice authors often experience anxiety and frustration while composing this very first sentence. Therefore, if this writing technique is introduced, it might help these inexperienced authors to have less anxiety and frustration and to feel more at ease when writing this sentence.

5. In general, a sentence should not be too short or too long because it might be inadequate or it might distract the reader's attention. This study found that opening sentences were composed of about 22 words. Therefore, it might be wise to take this length as an average length for opening sentences.

Suggestions for future research

This study examined only the first sentence of the introduction section of medical research articles, which is only the first element of the first move. Future research should investigate other inner elements, such as reporting previous findings that pave ways for elucidating gaps in research or giving reasons why a study was undertaken, the statement of the research gap as well as the statement of purpose.

Conclusion

The findings of this study suggest that opening sentences usually give facts, indicate importance, or express a combination of these two rhetorical functions. These sentences are also composed of about 22 words in the present simple in active voice. In addition, they are often a citation of previous findings.

Acknowledgements

The author thanks Assoc. Prof. Dr. Supat Sukamolisan, former Deputy Director for Research, and Asst. Prof. Dr. Kulaporn Hiranburana, former Deputy Director for Academic Affairs, for their help and moral support. Special thanks are extended to Jim Hutton, a language expert from the University of Edinburgh, who previewed and suggested improvement of the research proposal. Sincere appreciation is also extended to James Haft who always offers a helping hand.

The Author

Phan Banpho is an assistant professor and the Head of the Division of English for Science and Technology at CULI where he has been teaching for two decades. He is interested in applied linguistics as well as instructional technology. He has been contributing many articles to PASAA and Pasaparitat, journals published by CULI. He is the author of two books on writing computer programs using computer languages. He is also the author of two textbooks—*Reading Research Articles in Medical Sciences* and *Writing Research Articles in Medical Sciences*.

References

- Barber, C. L. (1985). Some measurable characteristics of modern scientific prose. In J. Swales (Ed). *Episodes in ESP* (pp. 3-16). Oxford: Pergamon Institute of English.
- Biber, D. et al. (1999). *Longman Grammar of Spoken and Written English*. Essex, UK: Longman.
- Brink, P. I., & Wood, M. J. (1994). *Basic Steps in Planning Nursing Research*. Boston: Jones and Bartlett Publishers.
- Hanania, E. A. S., & Karima, A. (1985). Verb form and rhetorical function in science writing: A study of MS theses in biology, chemistry, and physics. *The ESP Journal*, 4 (1), 49-58.

- Malcolm, L. (1987). What rules govern tense usage in scientific articles?
The ESP Journal, 6, 31-43.
- McCarthy, M. (1991). *Discourse analysis for language teachers*.
Cambridge: Cambridge University Press.
- McElroy, J. (1997). *Writing ahead: A process approach to academic writing*. Tokyo: Macmillan Language House.
- Oster, S. (1981). The use of tenses in "reporting past literature" in EST.
In L. Selinker, E. Tarone, & V. Hanzeli (Eds.). *English for academic and technical purposes: Studies in honor of Louis Trimble* (pp. 77-90). Rowley, MA: Newbury House.
- Smith, D. E. A. (1984). Medical discourse aspects of author's comment.
The ESP Journal 3, 25-36.
- Swales, J. (1984). Research into the structure of introductions to journal articles and its application to the teaching of academic writing. In R. Williams, J. Swales, and J. Kirkman (Eds.). *Common ground: Shared interests in ESP and communication studies* (pp. 77-86). Oxford: Pergamon.
- Swales, J. (1990). *Genre analysis*. Cambridge: Cambridge University Press.
- Tarone, E., Dwyer, E., Gillette, S., & Icke, V. (1985). On the use of the passive in two astrophysics journal papers. In J. Swales (Ed.). *Episodes in ESP* (pp. 191-205). Oxford: Pergamon Institute of English.
- Wingard, A. (1981). Some verb forms and functions in six medical texts. In L. Selinker, E. Tarone, & V. Hanzeli (Eds.). *English for academic and technical purposes: Studies in honor of Louis Trimble* (pp. 53-64). Rowley, MA: Newbury House Publishers.

